arXiv:0707.0902v2 [quant-ph] 18 Jul 2007

Flow Representation of the Bose–Hubbard Hamiltonian : General Case

Kazuyuki FUJII [*] and Tatsuo SUZUKI [†]

[*]Department of Mathematical Sciences
Yokohama City University
Yokohama, 236–0027
Japan

[†]Center for Educational Assistance
Shibaura Institute of Technology
Saitama, 337–8570
Japan

Abstract

In this paper the explicit flow representation to the Bose–Hubbard Hamiltonian is given in the general case. This representation may be useful in creating cat states for the system of atoms trapped in the optical ring.

To make macroscopic superpositions (a kind of cat states) of superfluid flows in a Bose–Einstein condensate (BEC) is really an interesting object in Quantum Mechanics. As a general introduction to BECs see for example [1] and [2]. When studying such states theoretically we usually use the Bose–Hubbard model, which is convenient enough for our purpose.

---

[*] E-mail address : fujii@yokohama-cu.ac.jp

[†] E-mail address : suzukita@aoni.waseda.jp ; i027110@sic.shibaura-it.ac.jp

We are especially interested in BECs trapped in an optical lattice in a ring geometry. On this subject there are many works, see [3], or [4], [5], [6] and their references.

In studying the model we often use a Fourier transform, which is a fundamental tool. In [4], [5], [6] the flow representation (a discrete Fourier transform) of the Bose–Hubbard Hamiltonian with three BECs is given to study a cat state. However, the system of three BECs is too small for our purpose.

We are studying a quantum computation based on Cavity QED (see [7] and [8]) and want to construct a hybrid system consisting of our method and some ideas (devices) coming from BECs, so it is important to know the flow representation of the Hamiltonian **in the general case**.

In this report we give the explicit flow representation to the Bose–Hubbard Hamiltonian in the general case. Our work may give a speed-up in studying the subject.

We treat in the paper the system of n harmonic oscillators $\{(a_1, a_1^\dagger), (a_2, a_2^\dagger), \cdots, (a_n, a_n^\dagger)\}$

$$[a_i, a_j^\dagger] = \delta_{ij}, \quad [a_i, a_j] = 0, \quad [a_i^\dagger, a_j^\dagger] = 0$$

trapped in an optical lattice in a ring geometry, see the following figure.



FIG : n atoms trapped in an optical lattice in a ring geometry

The Bose–Hubbard Hamiltonian in our context is

$$\mathrm{H} = -J\left(\sum_{i=1}^{n} a_{i+1}^\dagger a_i + h.c.\right) + \frac{U}{2}\sum_{i=1}^{n} {a_i^\dagger}^2 a_i^2 \tag{1}$$

where $a_{n+1} = a_1 \pmod{n}$ and J, U are some coupling constants consisting of the tunnelling strength, the interaction one respectively. See [3] or [6] and its references.

Let us rewrite the Hamiltonian by use of the matrix (vector) expression. We set

$$\mathbf{a} = (a_1, a_2, \cdots, a_{n-1}, a_n)^t$$

where t means the transpose and

$$\Sigma_1 = \begin{pmatrix} 0 & & & & & 1 \\ 1 & 0 & & & & \\ & 1 & 0 & & & \\ & & \ddots & \ddots & & \\ & & & 1 & 0 & \\ & & & & 1 & 0 \end{pmatrix}, \quad \Sigma_3 = \begin{pmatrix} 1 & & & & & \\ & \sigma & & & & \\ & & \sigma^2 & & & \\ & & & \ddots & & \\ & & & & \sigma^{n-2} & \\ & & & & & \sigma^{n-1} \end{pmatrix} \tag{2}$$

where $\sigma = \exp(2\pi\sqrt{-1}/n)$ which satisfies the relations

$$\sigma^n = 1, \quad \bar{\sigma} = \sigma^{n-1}, \quad 1 + \sigma + \cdots + \sigma^{n-1} = 0.$$

The generators $\{\Sigma_1, \Sigma_3\}$ are called the generalized Pauli matrices, which play an crucial role in qudit theory, see for example [9], [10], [11]. Then we have easily

$$\mathrm{H} = -J\left(\mathbf{a}^\dagger \Sigma_1 \mathbf{a} + h.c.\right) + \frac{U}{2}\sum_{i=1}^{n} {a_i^\dagger}^2 a_i^2. \tag{3}$$

What we want to do is to make the first term in (3) diagonal. For that let us remind the well–known decomposition

$$\Sigma_1 = W\Sigma_3 W^\dagger \;; \quad \Sigma_1^\dagger = W^\dagger \Sigma_3 W \quad (W^{-1} = W^\dagger) \tag{4}$$

with the generalized Walsh–Hadamard matrix (transformation) W defined by

$$W = \frac{1}{\sqrt{n}} \begin{pmatrix} 1 & 1 & 1 & \cdots & 1 & 1 \\ 1 & \sigma^{n-1} & \sigma^{2(n-1)} & \cdots & \sigma^{(n-2)(n-1)} & \sigma^{(n-1)^2} \\ 1 & \sigma^{n-2} & \sigma^{2(n-2)} & \cdots & \sigma^{(n-2)^2} & \sigma^{(n-1)(n-2)} \\ \vdots & \vdots & \vdots & & \vdots & \vdots \\ 1 & \sigma^{2} & \sigma^{4} & \cdots & \sigma^{2(n-2)} & \sigma^{2(n-1)} \\ 1 & \sigma & \sigma^{2} & \cdots & \sigma^{n-2} & \sigma^{n-1} \end{pmatrix}, \tag{5}$$

$$W^{\dagger} = \frac{1}{\sqrt{n}} \begin{pmatrix} 1 & 1 & 1 & \cdots & 1 & 1 \\ 1 & \sigma & \sigma^{2} & \cdots & \sigma^{n-2} & \sigma^{n-1} \\ 1 & \sigma^{2} & \sigma^{4} & \cdots & \sigma^{2(n-2)} & \sigma^{2(n-1)} \\ \vdots & \vdots & \vdots & & \vdots & \vdots \\ 1 & \sigma^{n-2} & \sigma^{2(n-2)} & \cdots & \sigma^{(n-2)^2} & \sigma^{(n-2)(n-1)} \\ 1 & \sigma^{n-1} & \sigma^{2(n-1)} & \cdots & \sigma^{(n-2)(n-1)} & \sigma^{(n-1)^2} \end{pmatrix}. \tag{6}$$

An interesting property of W used later is

$$W^2 = {W^{\dagger}}^2 = \begin{pmatrix} 1 & & & & & \\ & & & & & 1 \\ & & & & 1 & \\ & & & \iddots & & \\ & & 1 & & & \\ & 1 & & & & \end{pmatrix}. \tag{7}$$

It is worth noting that this property has been used in constructing the exchange gate (operator) in qudit theory (see [9] where $K = W^2$).

A comment may be in order. In the case of $n = 2$ we have

$$W = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}, \quad W^2 = \begin{pmatrix} 1 & \\ & 1 \end{pmatrix} = \mathbf{1}_2.$$

Surely, there is a (big) difference between $n = 2$ and $n \geq 3$.

Now we set

$$\alpha = W^{\dagger}\mathbf{a} \iff \mathbf{a} = W\alpha, \quad \alpha = (\alpha_1, \alpha_2, \cdots, \alpha_{n-1}, \alpha_n)^t, \tag{8}$$

or more explicitly

$$\alpha_1 = \frac{1}{\sqrt{n}}\sum_{i=1}^{n} a_i, \quad \alpha_2 = \frac{1}{\sqrt{n}}\sum_{i=1}^{n}\sigma^{i-1}a_i, \quad \cdots, \quad \alpha_n = \frac{1}{\sqrt{n}}\sum_{i=1}^{n}\sigma^{(i-1)(n-1)}a_i.$$

This change of "variables" is a discrete Fourier transform in quantum computation and is called the flow representation [1], see [4], [5], [6].

Under this transform the first term of (3) becomes

$$-2J\sum_{i=1}^{n}\cos\left(\frac{2\pi(i-1)}{n}\right)\alpha_i^{\dagger}\alpha_i, \tag{9}$$

which is just diagonal as required.

The real problem is to determine the second term of (3) in terms of (8). In fact, it is not so easy. The aim of this paper is to give the explicit form to the term and we obtain the following

Fundamental Formula

$$\sum_{i=1}^{n} a_i^{\dagger 2} a_i^2 = \frac{1}{n}\sum_{j=0}^{n-1}\left(\alpha^t W^2 \Sigma_1^j \alpha\right)^{\dagger}\left(\alpha^t W^2 \Sigma_1^j \alpha\right). \tag{10}$$

The proof will be given in [12]. It is interesting to note that W^2 in (7) is used in an essential manner.

Therefore the Bose–Hubbard Hamiltonian (1) becomes

$$\mathrm{H} = -2J\sum_{i=1}^{n}\cos\left(\frac{2\pi(i-1)}{n}\right)\alpha_i^{\dagger}\alpha_i + \frac{U}{2n}\sum_{j=0}^{n-1}\left(\alpha^t W^2 \Sigma_1^j \alpha\right)^{\dagger}\left(\alpha^t W^2 \Sigma_1^j \alpha\right) \tag{11}$$

in the **flow representation**.

Let us write down (11) explicitly for some special cases ($n = 3$, 4 and 5).

n=3 : ([4], [5], [6])

$$\begin{aligned}\mathrm{H} = &-J\left(2\alpha_1^{\dagger}\alpha_1 - \alpha_2^{\dagger}\alpha_2 - \alpha_3^{\dagger}\alpha_3\right) + \frac{U}{6}\left\{\left((\alpha_1^{\dagger})^2 + 2\alpha_2^{\dagger}\alpha_3^{\dagger}\right)\left(\alpha_1^2 + 2\alpha_2\alpha_3\right)\right. \\ &\left. + \left((\alpha_2^{\dagger})^2 + 2\alpha_1^{\dagger}\alpha_3^{\dagger}\right)\left(\alpha_2^2 + 2\alpha_1\alpha_3\right) + \left((\alpha_3^{\dagger})^2 + 2\alpha_1^{\dagger}\alpha_2^{\dagger}\right)\left(\alpha_3^2 + 2\alpha_1\alpha_2\right)\right\}.\end{aligned} \tag{12}$$

---

[1] we don't know whether this terminology is universal or not

n=4 :

$$\begin{aligned}\mathrm{H} = &-2J\left(\alpha_1^\dagger\alpha_1 - \alpha_3^\dagger\alpha_3\right) + \frac{U}{8}\left\{\left((\alpha_1^\dagger)^2 + 2\alpha_2^\dagger\alpha_4^\dagger + (\alpha_3^\dagger)^2\right)\left(\alpha_1^2 + 2\alpha_2\alpha_4 + \alpha_3^2\right)\right. \\ &+\left((\alpha_2^\dagger)^2 + 2\alpha_1^\dagger\alpha_3^\dagger + (\alpha_4^\dagger)^2\right)\left(\alpha_2^2 + 2\alpha_1\alpha_3 + \alpha_4^2\right) + 4\left(\alpha_1^\dagger\alpha_2^\dagger + \alpha_3^\dagger\alpha_4^\dagger\right)(\alpha_1\alpha_2 + \alpha_3\alpha_4) \\ &\left.+4\left(\alpha_1^\dagger\alpha_4^\dagger + \alpha_2^\dagger\alpha_3^\dagger\right)(\alpha_1\alpha_4 + \alpha_2\alpha_3)\right\}. \qquad (13)\end{aligned}$$

n=5 :

$$\begin{aligned}\mathrm{H} = &-\frac{J}{2}\left\{4\alpha_1^\dagger\alpha_1 + (\sqrt{5}-1)\left(\alpha_2^\dagger\alpha_2 + \alpha_5^\dagger\alpha_5\right) - (\sqrt{5}+1)\left(\alpha_3^\dagger\alpha_3 + \alpha_4^\dagger\alpha_4\right)\right\} \\ &+\frac{U}{10}\left\{\left((\alpha_1^\dagger)^2 + 2\alpha_2^\dagger\alpha_5^\dagger + 2\alpha_3^\dagger\alpha_4^\dagger\right)\left(\alpha_1^2 + 2\alpha_2\alpha_5 + 2\alpha_3\alpha_4\right) + \right. \\ &\left((\alpha_3^\dagger)^2 + 2\alpha_1^\dagger\alpha_5^\dagger + 2\alpha_2^\dagger\alpha_4^\dagger\right)\left(\alpha_3^2 + 2\alpha_1\alpha_5 + 2\alpha_2\alpha_4\right) + \\ &\left((\alpha_5^\dagger)^2 + 2\alpha_1^\dagger\alpha_4^\dagger + 2\alpha_2^\dagger\alpha_3^\dagger\right)\left(\alpha_5^2 + 2\alpha_1\alpha_4 + 2\alpha_2\alpha_3\right) + \\ &\left((\alpha_2^\dagger)^2 + 2\alpha_1^\dagger\alpha_3^\dagger + 2\alpha_4^\dagger\alpha_5^\dagger\right)\left(\alpha_2^2 + 2\alpha_1\alpha_3 + 2\alpha_4\alpha_5\right) + \\ &\left.\left((\alpha_4^\dagger)^2 + 2\alpha_1^\dagger\alpha_2^\dagger + 2\alpha_3^\dagger\alpha_5^\dagger\right)\left(\alpha_4^2 + 2\alpha_1\alpha_2 + 2\alpha_3\alpha_5\right)\right\}. \qquad (14)\end{aligned}$$

In this paper we treated the Bose–Hubbard Hamiltonian consisting of atoms trapped in the optical ring and gave the explicit flow representation (discrete Fourier transform) to it. This representation may be convenient in creating cat states for the system, which will be reported in another paper [12].

We conclude the paper by stating our motivation once more. We are studying a quantum computation (computer) based on Cavity QED, so to construct a more realistic model of (robust) quantum computer we have to study a hybrid system consisting of our method and some ideas (devices) coming from BECs. This is our future task.

References

[1] C. J. Pethick and H. Smith : Bose–Einstein Condensation in Dilute Gases, Cambridge University Press, Cambridge, 2001.

[2] L. P. Pitaevskii and S. Stringari : Bose–Einstein Condensation, Oxford University Press, Oxford, 2003.

[3] D. Jaksch, C. Bruder, J. I. Cirac, C. W. Gardiner and P. Zoller : Cold bosonic atoms in optical lattices, Phys. Rev. Lett, **81**, 3108 (1998).

[4] D. W. Hallwood, K. Burnett and J. Dunningham : Macroscopic superpositions of superfluid flows, New J. Phys, **8**, 180 (2006).

[5] J. Dunningham and D. Hallwood : Creation of macroscopic superpositions of flow states with Bose–Einstein condensates, Phys. Rev. A **74**, 023601 (2006).

[6] D. W. Hallwood, K. Burnett and J. Dunningham : The barriers to producing multiparticle superposition states in rotating Bose-Einstein condensates, to appear in The Journal of Modern Optics, quant-ph/0609077.

[7] K. Fujii, K. Higashida, R. Kato and Y. Wada : Cavity QED and Quantum Computation in the Weak Coupling Regime, J. Opt. B: Quantum and Semiclass. Opt, **6** (2004) 502, quant-ph/0407014.

[8] K. Fujii, K. Higashida, R. Kato and Y. Wada : Cavity QED and Quantum Computation in the Weak Coupling Regime II : Complete Construction of the Controlled–Controlled NOT Gate, Trends in Quantum Computing Research, Susan Shannon (Ed.), **Chapter 8**, Nova Science Publishers, 2006 and Computer Science and Quantum Computing, James E. Stones (Ed.), **Chapter 1**, Nova Science Publishers, 2007, quant-ph/0501046.

[9] K. Fujii : Exchange Gate on the Qudit Space and Fock Space, J. Opt. B : Quantum Semiclass. Opt, **5** (2003), S613, quant-ph/0207002.

[10] K. Fujii, K. Funahashi and T. Kobayashi : Jarlskog's Parametrization of Unitary Matrices and Qudit Theory, Int. J. Geom. Meth. Mod. Phys, **3** (2006), 269, quant-ph/0508006.

[11] K. Fujii : A New Algebraic Structure of Finite Quantum Systems and the Modified Bessel Functions, to appear in International Journal of Geometric Methods in Modern Physics, (vol.4, no.7), 2007, arXiv : 0704.1844 (quant-ph).

[12] K. Fujii and T. Suzuki : in preparation.